

With regard to birds it may be observed, that if spirit be injected down the windpipe, it will pass through almost the whole body by means of the air-cells. In the case of a quadruped preserved in spirit, or in the saline solution, it is proper to inject the preserving liquor into the abdominal cavity and intestinal canal.

BIBLIOGRAPHICAL NOTICES.

Supplement to the History of British Fishes. By William Yarrell, F.L.S., V.P.Z.S., Illustrated by Wood-cuts. London, Van Voorst, 8vo. 1839.

In an early number of the 'Magazine of Zoology and Botany,' we noticed the work to which that bearing the title above is a Supplement, considering it as one well executed in all its departments, and of much importance in bringing together our knowledge of a branch of the British Fauna which had not been investigated so minutely as either the zoology or ornithology; and moreover we looked upon it as a work which was sure to cause the production of much additional information. In the latter prophecy the Supplement before us bears ample testimony that we have not been deceived, while it also exhibits the zeal of the author in his interesting department, and the assiduity with which he is assisted, which is mainly to be attributed to the excitement given to our ichthyologists by the interest of his previous volumes.

The execution of the Supplement more than equals that of the former volumes. Thirty-one figures of fishes are given, besides vignettes illustrating structure or characters. Among the species the greater number are new to our Fauna, the others being improved representations of fishes previously illustrated, or taken from specimens where the former wood-cuts were, for want of better materials, copied from drawings or other engravings. In the letter-press several mistakes and inadvertencies have been corrected. We trust that Mr. Yarrell will continue to collect materials as he has already done, and we will venture to predict that it will not be long before they will again swell to an extent even beyond that which he has already presented to the public.

The following may be noticed as occurring among the more important corrections and additions to this Supplement:—Couch's *Serranus* Mr. Yarrell has found to be identical with the *Polyprion cernuum* of Cuv. and Val. An improved figure of the baagmaer or Deal fish is given from the representation of Professor Reinhart of

With regard to birds it may be observed, that if spirit be injected down the windpipe, it will pass through almost the whole body by means of the air-cells. In the case of a quadruped preserved in spirit, or in the saline solution, it is proper to inject the preserving liquor into the abdominal cavity and intestinal canal.

BIBLIOGRAPHICAL NOTICES.

Supplement to the History of British Fishes. By William Yarrell, F.L.S., V.P.Z.S., Illustrated by Wood-cuts. London, Van Voorst, 8vo. 1839.

In an early number of the 'Magazine of Zoology and Botany,' we noticed the work to which that bearing the title above is a Supplement, considering it as one well executed in all its departments, and of much importance in bringing together our knowledge of a branch of the British Fauna which had not been investigated so minutely as either the zoology or ornithology; and moreover we looked upon it as a work which was sure to cause the production of much additional information. In the latter prophecy the Supplement before us bears ample testimony that we have not been deceived, while it also exhibits the zeal of the author in his interesting department, and the assiduity with which he is assisted, which is mainly to be attributed to the excitement given to our ichthyologists by the interest of his previous volumes.

The execution of the Supplement more than equals that of the former volumes. Thirty-one figures of fishes are given, besides vignettes illustrating structure or characters. Among the species the greater number are new to our Fauna, the others being improved representations of fishes previously illustrated, or taken from specimens where the former wood-cuts were, for want of better materials, copied from drawings or other engravings. In the letter-press several mistakes and inadvertencies have been corrected. We trust that Mr. Yarrell will continue to collect materials as he has already done, and we will venture to predict that it will not be long before they will again swell to an extent even beyond that which he has already presented to the public.

The following may be noticed as occurring among the more important corrections and additions to this Supplement:—Couch's *Serranus* Mr. Yarrell has found to be identical with the *Polyprion cernuum* of Cuv. and Val. An improved figure of the baagmaer or Deal fish is given from the representation of Professor Reinhart of

With regard to birds it may be observed, that if spirit be injected down the windpipe, it will pass through almost the whole body by means of the air-cells. In the case of a quadruped preserved in spirit, or in the saline solution, it is proper to inject the preserving liquor into the abdominal cavity and intestinal canal.

BIBLIOGRAPHICAL NOTICES.

Supplement to the History of British Fishes. By William Yarrell, F.L.S., V.P.Z.S., Illustrated by Wood-cuts. London, Van Voorst, 8vo. 1839.

In an early number of the 'Magazine of Zoology and Botany,' we noticed the work to which that bearing the title above is a Supplement, considering it as one well executed in all its departments, and of much importance in bringing together our knowledge of a branch of the British Fauna which had not been investigated so minutely as either the zoology or ornithology; and moreover we looked upon it as a work which was sure to cause the production of much additional information. In the latter prophecy the Supplement before us bears ample testimony that we have not been deceived, while it also exhibits the zeal of the author in his interesting department, and the assiduity with which he is assisted, which is mainly to be attributed to the excitement given to our ichthyologists by the interest of his previous volumes.

The execution of the Supplement more than equals that of the former volumes. Thirty-one figures of fishes are given, besides vignettes illustrating structure or characters. Among the species the greater number are new to our Fauna, the others being improved representations of fishes previously illustrated, or taken from specimens where the former wood-cuts were, for want of better materials, copied from drawings or other engravings. In the letter-press several mistakes and inadvertencies have been corrected. We trust that Mr. Yarrell will continue to collect materials as he has already done, and we will venture to predict that it will not be long before they will again swell to an extent even beyond that which he has already presented to the public.

The following may be noticed as occurring among the more important corrections and additions to this Supplement:—Couch's *Serranus* Mr. Yarrell has found to be identical with the *Polyprion cernuum* of Cuv. and Val. An improved figure of the baagmaer or Deal fish is given from the representation of Professor Reinhart of

With regard to birds it may be observed, that if spirit be injected down the windpipe, it will pass through almost the whole body by means of the air-cells. In the case of a quadruped preserved in spirit, or in the saline solution, it is proper to inject the preserving liquor into the abdominal cavity and intestinal canal.

BIBLIOGRAPHICAL NOTICES.

Supplement to the History of British Fishes. By William Yarrell, F.L.S., V.P.Z.S., Illustrated by Wood-cuts. London, Van Voorst, 8vo. 1839.

In an early number of the 'Magazine of Zoology and Botany,' we noticed the work to which that bearing the title above is a Supplement, considering it as one well executed in all its departments, and of much importance in bringing together our knowledge of a branch of the British Fauna which had not been investigated so minutely as either the zoology or ornithology; and moreover we looked upon it as a work which was sure to cause the production of much additional information. In the latter prophecy the Supplement before us bears ample testimony that we have not been deceived, while it also exhibits the zeal of the author in his interesting department, and the assiduity with which he is assisted, which is mainly to be attributed to the excitement given to our ichthyologists by the interest of his previous volumes.

The execution of the Supplement more than equals that of the former volumes. Thirty-one figures of fishes are given, besides vignettes illustrating structure or characters. Among the species the greater number are new to our Fauna, the others being improved representations of fishes previously illustrated, or taken from specimens where the former wood-cuts were, for want of better materials, copied from drawings or other engravings. In the letter-press several mistakes and inadvertencies have been corrected. We trust that Mr. Yarrell will continue to collect materials as he has already done, and we will venture to predict that it will not be long before they will again swell to an extent even beyond that which he has already presented to the public.

The following may be noticed as occurring among the more important corrections and additions to this Supplement:—Couch's *Serranus* Mr. Yarrell has found to be identical with the *Polyprion cernuum* of Cuv. and Val. An improved figure of the baagmaer or Deal fish is given from the representation of Professor Reinhart of

Copenhagen, together with extracts from that naturalist's not easily accessible memoir. There is a beautifully cut figure of *Crenilabrus rupestris*, which formed the subject of an interesting paper by Mr. Selby in an early number of this periodical, and which (since attention has been directed to the species) has been met with on several parts of the English and Irish shores. In the paper alluded to 'Mag. of Zool. and Bot.' i. p. 137, the third specimen is said to have been found on the shore near "*Barncleugh*:" there is no such place, the name is a misprint for *Bamborough*, which we regret to see copied by Mr. Yarrell, as well as by foreign writers without blame on their parts, and we trust that this notice will tend to correction. The *Pomeranian* Bream, *C. Buggenhaggi*, is a fine addition to our *Cyprinidæ*, and has been taken in Dagenhambreach, Essex, and by Mr. Thompson in the river Logan near Belfast. Of the very singular genus *Hemiramphus*, a fish is represented under the title of *H. Europæus*, detected several years since by Mr. Couch of Polpero. In the description Mr. Yarrell remarks, "one question may be hazarded:—Is this fish, with its unequally developed jaws, the very young state of our common Garfish, *Belone vulgaris*?" and we would request our ichthyologists to attend to this query, very interesting and important, whether it proves us to have a British member of the genus *Hemiramphus*, or that the lengthened jaw is only a peculiar structure in another division; and perhaps touching upon another question immediately suggested by Mr. Yarrell's surmise,—whether or not the *Hemiramphi* may not be all a young or immature form of this fish, and if so what is the use of the elongation of the under jaw at this early period? Among the *Salmonidæ* we have a second figure of *S. ferox* from an Irish specimen, authenticating its Irish range. Mr. Yarrell also mentions having received specimens of trout weighing 32 and 34 lbs. from Lake Wenern, and which are considered identical with the British fish. A species of smelt, *Osmerus Hebridicus*, taken near the Isle of Bute, is given as new to Britain and to ichthyology. The *powan* of Loch Lomond, and *pollan* of Loch Neagh, are also now figured. *Platessa elongata*, taken at Stoford, in Bridgewater Bay, is described as new to ichthyology. Among the British Sturgeons ichthyologists will find interesting employment, and we refer to Mr. Yarrell's finely cut figures of the heads of two, and to his short observations for excitement. *Echinorhinus spinosus* is a very interesting addition, but surely there is much difference between Mr. Yarrell's figure and Dr. Smith's in the proportion and relative positions of the members, and also in the form of the teeth. We judge now only from the figures before us.

Copenhagen, together with extracts from that naturalist's not easily accessible memoir. There is a beautifully cut figure of *Crenilabrus rupestris*, which formed the subject of an interesting paper by Mr. Selby in an early number of this periodical, and which (since attention has been directed to the species) has been met with on several parts of the English and Irish shores. In the paper alluded to 'Mag. of Zool. and Bot.' i. p. 137, the third specimen is said to have been found on the shore near "*Barncleugh*:" there is no such place, the name is a misprint for *Bamborough*, which we regret to see copied by Mr. Yarrell, as well as by foreign writers without blame on their parts, and we trust that this notice will tend to correction. The *Pomeranian* Bream, *C. Buggenhaggi*, is a fine addition to our *Cyprinidæ*, and has been taken in Dagenhambreach, Essex, and by Mr. Thompson in the river Logan near Belfast. Of the very singular genus *Hemiramphus*, a fish is represented under the title of *H. Europæus*, detected several years since by Mr. Couch of Polpero. In the description Mr. Yarrell remarks, "one question may be hazarded:—Is this fish, with its unequally developed jaws, the very young state of our common Garfish, *Belone vulgaris*?" and we would request our ichthyologists to attend to this query, very interesting and important, whether it proves us to have a British member of the genus *Hemiramphus*, or that the lengthened jaw is only a peculiar structure in another division; and perhaps touching upon another question immediately suggested by Mr. Yarrell's surmise,—whether or not the *Hemiramphi* may not be all a young or immature form of this fish, and if so what is the use of the elongation of the under jaw at this early period? Among the *Salmonidæ* we have a second figure of *S. ferox* from an Irish specimen, authenticating its Irish range. Mr. Yarrell also mentions having received specimens of trout weighing 32 and 34 lbs. from Lake Wenern, and which are considered identical with the British fish. A species of smelt, *Osmerus Hebridicus*, taken near the Isle of Bute, is given as new to Britain and to ichthyology. The *powan* of Loch Lomond, and *pollan* of Loch Neagh, are also now figured. *Platessa elongata*, taken at Stoford, in Bridgewater Bay, is described as new to ichthyology. Among the British Sturgeons ichthyologists will find interesting employment, and we refer to Mr. Yarrell's finely cut figures of the heads of two, and to his short observations for excitement. *Echinorhinus spinosus* is a very interesting addition, but surely there is much difference between Mr. Yarrell's figure and Dr. Smith's in the proportion and relative positions of the members, and also in the form of the teeth. We judge now only from the figures before us.

Copenhagen, together with extracts from that naturalist's not easily accessible memoir. There is a beautifully cut figure of *Crenilabrus rupestris*, which formed the subject of an interesting paper by Mr. Selby in an early number of this periodical, and which (since attention has been directed to the species) has been met with on several parts of the English and Irish shores. In the paper alluded to 'Mag. of Zool. and Bot.' i. p. 137, the third specimen is said to have been found on the shore near "*Barncleugh*:" there is no such place, the name is a misprint for *Bamborough*, which we regret to see copied by Mr. Yarrell, as well as by foreign writers without blame on their parts, and we trust that this notice will tend to correction. The *Pomeranian* Bream, *C. Buggenhaggi*, is a fine addition to our *Cyprinidæ*, and has been taken in Dagenhambreach, Essex, and by Mr. Thompson in the river Logan near Belfast. Of the very singular genus *Hemiramphus*, a fish is represented under the title of *H. Europæus*, detected several years since by Mr. Couch of Polpero. In the description Mr. Yarrell remarks, "one question may be hazarded:—Is this fish, with its unequally developed jaws, the very young state of our common Garfish, *Belone vulgaris*?" and we would request our ichthyologists to attend to this query, very interesting and important, whether it proves us to have a British member of the genus *Hemiramphus*, or that the lengthened jaw is only a peculiar structure in another division; and perhaps touching upon another question immediately suggested by Mr. Yarrell's surmise,—whether or not the *Hemiramphi* may not be all a young or immature form of this fish, and if so what is the use of the elongation of the under jaw at this early period? Among the *Salmonidæ* we have a second figure of *S. ferox* from an Irish specimen, authenticating its Irish range. Mr. Yarrell also mentions having received specimens of trout weighing 32 and 34 lbs. from Lake Wenern, and which are considered identical with the British fish. A species of smelt, *Osmerus Hebridicus*, taken near the Isle of Bute, is given as new to Britain and to ichthyology. The *powan* of Loch Lomond, and *pollan* of Loch Neagh, are also now figured. *Platessa elongata*, taken at Stoford, in Bridgewater Bay, is described as new to ichthyology. Among the British Sturgeons ichthyologists will find interesting employment, and we refer to Mr. Yarrell's finely cut figures of the heads of two, and to his short observations for excitement. *Echinorhinus spinosus* is a very interesting addition, but surely there is much difference between Mr. Yarrell's figure and Dr. Smith's in the proportion and relative positions of the members, and also in the form of the teeth. We judge now only from the figures before us.

Copenhagen, together with extracts from that naturalist's not easily accessible memoir. There is a beautifully cut figure of *Crenilabrus rupestris*, which formed the subject of an interesting paper by Mr. Selby in an early number of this periodical, and which (since attention has been directed to the species) has been met with on several parts of the English and Irish shores. In the paper alluded to 'Mag. of Zool. and Bot.' i. p. 137, the third specimen is said to have been found on the shore near "*Barncleugh*:" there is no such place, the name is a misprint for *Bamborough*, which we regret to see copied by Mr. Yarrell, as well as by foreign writers without blame on their parts, and we trust that this notice will tend to correction. The *Pomeranian* Bream, *C. Buggenhaggi*, is a fine addition to our *Cyprinidæ*, and has been taken in Dagenhambreach, Essex, and by Mr. Thompson in the river Logan near Belfast. Of the very singular genus *Hemiramphus*, a fish is represented under the title of *H. Europæus*, detected several years since by Mr. Couch of Polpero. In the description Mr. Yarrell remarks, "one question may be hazarded:—Is this fish, with its unequally developed jaws, the very young state of our common Garfish, *Belone vulgaris*?" and we would request our ichthyologists to attend to this query, very interesting and important, whether it proves us to have a British member of the genus *Hemiramphus*, or that the lengthened jaw is only a peculiar structure in another division; and perhaps touching upon another question immediately suggested by Mr. Yarrell's surmise,—whether or not the *Hemiramphi* may not be all a young or immature form of this fish, and if so what is the use of the elongation of the under jaw at this early period? Among the *Salmonidæ* we have a second figure of *S. ferox* from an Irish specimen, authenticating its Irish range. Mr. Yarrell also mentions having received specimens of trout weighing 32 and 34 lbs. from Lake Wenern, and which are considered identical with the British fish. A species of smelt, *Osmerus Hebridicus*, taken near the Isle of Bute, is given as new to Britain and to ichthyology. The *powan* of Loch Lomond, and *pollan* of Loch Neagh, are also now figured. *Platessa elongata*, taken at Stoford, in Bridgewater Bay, is described as new to ichthyology. Among the British Sturgeons ichthyologists will find interesting employment, and we refer to Mr. Yarrell's finely cut figures of the heads of two, and to his short observations for excitement. *Echinorhinus spinosus* is a very interesting addition, but surely there is much difference between Mr. Yarrell's figure and Dr. Smith's in the proportion and relative positions of the members, and also in the form of the teeth. We judge now only from the figures before us.